

Формализм Ходжкина-Хаксли

«нейробиофизика», лекция 2


А. Р. Браже

- <http://neurondynamics.epfl.ch/>
- <http://neurondynamics-exercises.readthedocs.io/en/latest/>

Neuronal Dynamics

From single neurons to networks and models of cognition

Wulfram Gerstner, Werner M. Kistler, Richard Naud and Liam Paninski



What happens in our brain when we make a decision? What triggers a neuron to send out a signal? What is the neural code? This textbook for advanced undergraduate and beginning graduate students provides a thorough and up-to-date introduction to the fields of computational and theoretical neuroscience.


 [Read the online version of the book »](#)

 [Order the book »](#)


 [First-edition Errata »](#)

 [Python Exercises »](#)


Excitability, spiking and bursting




Neural Excitability



Periodic Spiking






Periodic Bursting




Prescott SA, De Koninck Y, Sejnowski TJ (2008) *PLoS Comput Biol* 4(10): e1000198. doi:10.1371/journal.pcbi.1000198


Ion currents in the squid giant axon




Ходжкин-Хаксли: разделение токов на I_{Na} и I_K




B Currents from large depolarization



A Currents from small depolarization



K⁺ and Na⁺ currents




Замена ионного состава среды

Использование блокаторов



Potassium current in the HH model

Dependence of K conductance on potential







$$n_{\infty} = \frac{\alpha_n}{\alpha_n + \beta_n}$$

$$\tau_n = \frac{1}{\alpha_n + \beta_n}$$




Sodium current in the HH model

$$I_{Na} = \bar{g}_{Na} m^3 h (V - E_{Na})$$



The Hodgkin-Huxley formalism



Membrane potential dynamics is governed by an ODE:

$$C_m \frac{dV}{dt} = I_{\text{electrode}} - \bar{g}_L (V - E_L) - \bar{g}_{Na} m^3 h (V - E_{Na}) - \bar{g}_K n^4 (V - E_K)$$

General summary

- 1) $I_i = g_i (V - E_{iNernst})$ Instantaneous I-V relationship is linear
- 2) $g_i = \bar{g}_i w^\gamma v^\delta$ Instantaneous conductance is maximal conductance weighted by fraction of open channels (2 types of gates)
- 3) $\frac{dw}{dt} = \frac{1}{\tau_w} (w_\infty - w) \equiv \alpha_w (1 - w) - \beta_w w$ Linear gate kinetics
- 4) $\alpha_w = f_1(V), \beta_w = f_2(V)$ Gate kinetic rates are nonlinear functions of V_m

$$I_K = \bar{g}_K n^4 (V - E_K)$$

K-current (through Kv channels)

$$I_{Na} = \bar{g}_{Na} m^3 h (V - E_{Na})$$


Na-current (through Nv-channels)

$$I_L = \bar{g}_L (V - E_L)$$

Non potential-sensitive leak

$$\frac{dn}{dt} = \alpha_n (1 - n) - \beta_n n$$

The Hodgkin-Huxley formalism




Membrane potential dynamics is governed by an ODE:

$$C_m \frac{dV}{dt} = I_{electrode} - \bar{g}_L (V - E_L) - \bar{g}_{Na} m^3 h (V - E_{Na}) - \bar{g}_K n^4 (V - E_K)$$

General summary

- 1) $I_i = g_i (V - E_{iNernst})$ Instantaneous I-V relationship is linear
- 2) $g_i = \bar{g}_i w^\gamma v^\delta$ Instantaneous conductance is maximal conductance weighted by fraction of open channels (2 types of gates)
- 3) $\frac{dw}{dt} = \frac{1}{\tau_w} (w_\infty - w) \equiv \alpha_w (1 - w) - \beta_w w$ Linear gate kinetics
- 4) $\alpha_w = f_1(V), \beta_w = f_2(V)$ Gate kinetic rates are nonlinear functions of V_m



α_n


β_n

80

18



10

The Hodgkin-Huxley formalism




Membrane potential dynamics is governed by an ODE:

$$C_m \frac{dV}{dt} = I_{\text{electrode}} - \bar{g}_L(V - E_L) - \bar{g}_{Na} m^3 h (V - E_{Na}) - \bar{g}_K n^4 (V - E_K)$$




Effect of temperature


$$\alpha(T) = \alpha(T_0) Q_{10}^{\frac{T-T_0}{10}}$$






Response of the HH model to current pulses: Class 2




Generation and conduction of spike in the squid axon



$$C_m \frac{dV}{dt} = I_{electrode} - \bar{g}_L(V - E_L) - \bar{g}_{Na} m^3 h (V - E_{Na}) - \bar{g}_K n^4 (V - E_K)$$



The zoo of ion currents




$I_{Na,p} + I_K$ -model:

$$C\dot{V} = I - \bar{g}_K n(V - E_K) - \bar{g}_{Na} m_\infty(V)(V - E_{Na}) - g_l(V - E_l)$$

$$\tau_n \dot{n} = (n_\infty(V) - n)$$


$$x_\infty = \frac{1}{1 + \exp(\frac{V_x^{0.5} - V}{k_x})}$$








Nullclines:

$$\dot{V} = 0 \rightarrow n(V) = \frac{I - \bar{g}_{Na} m_\infty(V - E_{Na}) - g_l(V - E_l)}{\bar{g}_k(V - E_k)}$$

$$\dot{n} = 0 \rightarrow n(V) = n_\infty(V)$$




More detailed channel models




Markov models and kinetic schemes

Na_v-channel (Vandenberg & Bezanilla 1991):



Why does everyone still uses HH formalism and independent gating particles?


- less parameters to fit
- less equations
- accurate enough
- when sub-ms accuracy is not required
- for large ensembles of channels



$$k_\mu = \frac{k_B T}{h} \exp\left(-\frac{\Delta G_\mu}{RT}\right) = \frac{k_B T}{h} \exp\left(\frac{\Delta S_\mu}{R}\right) \exp\left(-\frac{\Delta H_\mu}{RT}\right)$$

$$\Delta H_\mu(V) = \Delta H_\mu^{(0)} - \delta_\mu z_\mu F V,$$

$$\Delta H_{-\mu}(V) = \Delta H_{-\mu}^{(0)} + (1 - \delta_\mu) z_\mu F V.$$







Thermodynamic-based models

$$x_{\infty} = \frac{1}{1 + \exp(-(V - V_{1/2})/\sigma)},$$


$$\tau_x = \frac{1}{\alpha'(V) + \beta'(V)} + \tau_0,$$

$$\alpha'_x(V) = K \exp\left(\frac{\delta(V - V_{1/2})}{\sigma}\right)$$


$$\beta'_x(V) = K \exp\left(\frac{-(1 - \delta)(V - V_{1/2})}{\sigma}\right).$$




Excitability, spiking and bursting




Neural Excitability



Periodic Spiking



Periodic Bursting



Prescott SA, De Koninck Y, Sejnowski TJ (2008) *PLoS Comput Biol* 4(10): e1000198. doi:10.1371/journal.pcbi.1000198


Instantaneous persistent sodium and potassium current model ($I_{Na,p} + I_K$)

System:

$$C\dot{V} = I - \bar{g}_K n(V - E_K) - \bar{g}_{Na} m_\infty(V)(V - E_{Na}) - g_l(V - E_l)$$

$$\tau_n \dot{n} = (n_\infty(V) - n)$$

$$x_\infty = \frac{1}{1 + \exp(\frac{V_x^{0.5} - V}{k_x})}$$



Nullclines:

$$\dot{V} = 0 \rightarrow n(V) = \frac{I - \bar{g}_{Na} m_\infty(V - E_{Na}) - g_l(V - E_l)}{\bar{g}_K(V - E_K)}$$

$$\dot{n} = 0 \rightarrow n(V) = n_\infty(V)$$

